

Continuing professional development activity for <u>dietitians</u>

ACTIVITY 69 You can obtain 3 CEU's for reading the article: "PRIMARY HEALTH CARE FACILITY

SAJCN CPD activity No 68 and 69 - 2010

ACTIVITY 68

You can obtain 3 CEU's for reading the article: "NUTRITIONAL ASSESSMENT OF THE CRITICALLY ILL PATIENT" and answering the accompanying questions. This article has been accredited for CEU's for dietitians. Ref number: DT/A01/P00008/2010/00001					
PLEASE ANSWER ALL THE QUESTIONS (True or False)					
1.	About $70-80\%$ of patients are under- or malnourished upon admission to hospitals.	☐ True ☐ False			
2.	Lack of early nutritional assessment and treatment plays a role in weight loss during hospitalization.	True True			
3.	Elderly patients with chronic obstructive pulmonary disease and diabetes have a significantly poorer nutritional status on admission and a high readmission rate.	True			
4.	Utilizing the Harris Benedict formula without a stress factors for patients who suffered a central venous incident, provides sufficient nutrition support.	☐ True ☐ False			
5.	The hallmark of the refeeding syndrome is hypophosphataemia.	☐ True ☐ False			
6.	Current screening tools can be utilized to identify at risk critically ill obese patients.	True False			
7.	The NRS-2002 screening tool predicts incidence and severity of complications in GIT surgery patients better than the NRS and the NRI screening tools.	True False			
8.	Assessment of ingestion, deglutition and energy intake will help to determine the need for consistency adaptations and/	True False			

INFRASTRUCTURE AND SERVICES AND THE NUTRITIONAL STATUS OF CHILDREN 0 TO 71 MONTHS OLD AND THEIR CAREGIVERS ATTENDING THESE FACILITIES IN FOUR RURAL DISTRICTS IN THE EASTERN CAPE AND KWAZULU- NATAL PROVINCES, SOUTH AFRICA" and answering the accompanying questions. This article has been accredited for CEU's for dietitians. Ref number: DT/A01/P00008/2010/00002					
PLEASE ANSWER ALL THE QUESTIONS (True or False)					
1.	In this study, the researchers used different questionnaires for the Primary Health Care (PHC) facility nurses and the caregivers.	☐ True ☐ False			
2.	The researchers did not obtain permission to enter the health facilities as no invasive procedures were performed in the study.	☐ True ☐ False			
3.	In this study, 80% of the PHC facilities in the Eastern Cape (EC) and 75% in KuaZulu-natal (KZN) had access to safe drinking water.	☐ True ☐ False			
4.	All the PHC facilities that were included in the study had malnutrition registers.	☐ True ☐ False			
5.	This study showed that caregivers in rural districts of the EC and KZN do not visit facilities other than the local PHC facility for health care.	True True			
6.	The caregivers reported diarrhoea as the leading cause of under-five child deaths.	☐ True ☐ False			
7.	This study showed that the prevalence of stunting in 6 to 59-month-old children in rural districts of the EC and KZN was high according to WHO classification.	True True			
8.	The study also showed that the prevalence of wasting in 6 to 59-month-old children in rural districts of the EC was medium according to WHO classification.	☐ True ☐ False			
9.	This study showed that infrastructural problems, insufficient resources and the existence of cultural practices were among the key factors influencing the quality of health service delivery.	True True False			
10.	According to the literature, nurses interpret nutritional status differently than dietitians because of differences in training.	☐ True ☐ False			
11.	Homebirths in rural districts of the EC and KZN are an ideal option to promote safe maternal and child health practices.	True True			

12.	The authors argued that the shortcomings in infrastructure, services, human resources and nurses' training must be addressed before a difference in the health and nutritional status of vulnerable groups in South Africa can be expected.	True False
-----	---	---------------

HOW TO EARN YOUR CEUS

complications.

or supplementary foods/enteral products.

volumes are 200 - 250ml.

status in critically ill patients.

9. The most recent recommendations for gastric residual

10. The serum proteins, retinol binding protein and pre-albumin,

deficiency and therefore they are good indicators of protein

11. The NICE-sugar study showed an increased 90-day mortality

percentile has been reported to predict mortality and major

with strict blood glucose control in critically ill patients.

12. Upper mid-arm circumference (UMAC) below the 15th

are not affected by the acute phase response or zinc

1. Please check your name and HPCSA number. 2. Read the article and answer all the questions. 3. Only ONE answer sheet may be submitted per article. 4. Indicate your answers to the questions by coloring the appropriate block on this answer sheet. 5. This CPD questionnaire can now be completed online at <u>www.sajcn.co.za</u>. Click on "CPD" on the main page. You are now linked to the ecpdsa website. Register with the ecpdsa website, log in and choose the SAJCN CPD activity for this issue. Please provide your fax number and e-mail address for your processed certificate. 6. Only torn-out copies will be accepted by post. Please sign the document and post it to Postnet Suite 55, Private Bagx22, Tygervalley, Cape Town, 7536. 7. Answer sheets must be submitted **not later than March 2010**. Answer sheets received after this date will not be processed. 8. You will earn 3 CEUs per article if you answer more than 70% of the questions correctly. A score of less than 70% will not earn you any CEUs. 9. Make a photocopy for your own records in case your form is lost in the mail.

True

False

True

True

True

False

False

False

HPCSA number: DT	
Surname as registered with HPCSA:	Initials:
Cell phone number:	Fax No.:
E-mail address*:	[*NB: please supply e-mail address (if available) for certificate delivery – please write clearly]
Postal address:	
Code:	Signature
Full member of: ADSA: 🗌 yes 🗌 no 🛛 NSSA: 🗌 yes 🥅 no	